



علاقات متعددات حدود بوبكر بصيغة تسلسل قوالب Fermat

كارم بوبكر¹, لين زهانغ²

قسم الفيزياء والكيمياء¹, المدرسة العليا للعلوم والتكنولوجيات بتونس (م.ع.ع.ت.ت.)

جامعة تونس، الجمهورية التونسية

قسم الرياضيات², جامعة زهو جيانغ, هانغ زهو

جمهورية الصين الشعبية 310027

الملخص:

في هذا البحث تمت دراسة متعددات حدود بوبكر (Boubaker polynomials). باستخدام تحليل قوالب ريوordan (Riordan) أمكن إيجاد علاقات حدود بوبكر بمتعددات حدود شيبيشيف (Chebyshev) من النوع الأول والثاني وكذلك متعددات حدود فيرمات (Fermat).

تم إستنتاج بعض العلاقات التي يمكن الإستفادة منها في التطبيقات الفيزيائية المتعلقة بمثال التفصيل التحليلي لمتعددات حدود بوبكر .(Boubaker Polynomials Expansion Scheme: BPES)



University of Bahrain

Journal of the Association of Arab Universities for Basic and Applied Sciences

www.elsevier.com/locate/jaaubas
www.sciencedirect.com



ORIGINAL ARTICLE

Fermat-linked relations for the Boubaker polynomial sequences via Riordan matrices analysis

Karem Boubaker ^a, Lin Zhang ^{b,*}

^a ESSTT/ 63 Rue Sidi Jabeur 5100, Mahdia, Tunisia

^b Institute of Mathematics, Hangzhou Dianzi University, Hangzhou 310037, PR China

Available online 2 July 2012

KEYWORDS

Riordan matrices;
Boubaker polynomials

Abstract The Boubaker polynomials are investigated in this paper. Using a Riordan matrices analysis, a sequence of relations outlining the relations with Chebyshev and Fermat polynomials has been obtained. The obtained expressions are a meaningful supplement to recent applied physics studies using the Boubaker polynomials expansion scheme (BPES).

© 2012 University of Bahrain. Production and hosting by Elsevier B.V. All rights reserved.

1. Introduction

Polynomial expansion methods are extensively used in many mathematical and engineering fields to yield meaningful results for both numerical and analytical analysis (Alvarez et al., 2005; Bender and Dunne, 1988; Choi et al., 2004; Guertz et al., 2000; Koelink, 1994; Okada et al., 2006; Philippou and Georghiou, 1989; Sloan and Womersley, 2002). Among the most frequently used polynomials, the Boubaker polynomials are one of the interesting tools which were associated to several applied physics problems as well as the related polynomials such as the Boubaker–Turki polynomials (Boubaker, 2007, 2008; Bagula and Adamson, 2008; Sloane, 2008; Ghanouchi

et al., 2008; Slama et al., 2008b,c; Ghrib et al., 2008), the 4q Boubaker polynomials (Zhao et al., 2009) and the Boubaker–Zhao polynomials (Zhao et al., xxxx). For example, for some resolution purposes, a function $f(r)$ is expressed as an infinite nonlinear expansion of Boubaker–Zhao polynomials

$$f(r) = \lim_{N \rightarrow +\infty} \left[\frac{1}{2N} \sum_{n=1}^N \zeta_n \widehat{B}_{4n} \left(r \frac{\alpha_n}{R} \right) \right], \quad (1.1)$$

where α_n are the minimal positive roots of the Boubaker 4n-order polynomials \widehat{B}_{4n} , R is a maximum radial range and ζ_n are coefficients to be determined using the expression of $f(r)$. Since the Boubaker 4n-order polynomials have the particular properties: for any n ,

$$\begin{cases} \widehat{B}_{4n}(r)|_{r=0} = -2, \\ \frac{\partial \widehat{B}_{4n}(r)}{\partial r} = 0, \\ \frac{\partial^2 \widehat{B}_{4n}(r)}{\partial r^2} = 4n(n-1). \end{cases} \quad (1.2)$$

The related system (1.3) is induced:

$$\begin{cases} f(0) = \lim_{N \rightarrow +\infty} \left[\frac{1}{2N} \sum_{n=1}^N \zeta_n \widehat{B}_{4n} \left(r \frac{\alpha_n}{R} \right) \right] |_{r=0} = -\frac{1}{N} \sum_{n=1}^N \zeta_n, \\ f(R) = \lim_{N \rightarrow +\infty} \left[\frac{1}{2N} \sum_{n=1}^N \zeta_n \widehat{B}_{4n} \left(r \frac{\alpha_n}{R} \right) \right] |_{r=R} = 0, \\ \frac{\partial f(r)}{\partial r} |_{r=0} = \lim_{N \rightarrow +\infty} \left[\frac{1}{2N} \sum_{n=1}^N \zeta_n \frac{\partial (\widehat{B}_{4n} \left(r \frac{\alpha_n}{R} \right))}{\partial r} \right] |_{r=0} = 0. \end{cases} \quad (1.3)$$

* Corresponding author.

E-mail addresses: mmbb1112000@yahoo.fr (K. Boubaker), god_yalin@163.com (L. Zhang).

1815-3852 © 2012 University of Bahrain. Production and hosting by Elsevier B.V. All rights reserved.

Peer review under responsibility of University of Bahrain.
<http://dx.doi.org/10.1016/j.jaubas.2012.05.004>



2. The Boubaker polynomials

The first monomial definition of the Boubaker polynomials (Awojoyogbe and Boubaker, 2009; Boubaker, 2007; Chaouchi et al., 2007; Labiad and Boubaker, 2007) appeared in a physical study that yielded an analytical solution to the heat equation inside a physical model (Labiadh et al., 2008; Slama et al., 2008a). This monomial definition is expressed by (2.1):

Definition 2.1. A monomial definition of the Boubaker polynomials is:

$$B_n(X) \stackrel{\text{def}}{=} \sum_{p=0}^{\xi(n)} \left[\frac{n-4p}{n-p} \binom{p}{n-p} \right] (-1)^p X^{n-2p}, \quad (2.1)$$

where $\xi(n) = \lfloor \frac{n}{2} \rfloor \stackrel{\text{def}}{=} \frac{2n+(-1)^n-1}{4}$ (The symbol $\lfloor * \rfloor$ designates the floor function). Their coefficients could be defined through a recursive formula (2.2):

$$\begin{cases} B_n(X) = \sum_{j=0}^{\xi(n)} [b_{nj} X^{n-2j}], \\ b_{n,0} = 1, \\ b_{n,1} = -(n-4), \\ b_{n,j+1} = \frac{(n-2j)(n-2j-1)}{(j+1)(n-j-1)} \cdot \frac{n-4j-4}{n-4j} \cdot b_{nj}, \\ b_{n,\xi(n)} = \begin{cases} (-1)^{\frac{n}{2}} \cdot 2 & \text{if } n \text{ even,} \\ (-1)^{\frac{n+1}{2}} \cdot (n-2) & \text{if } n \text{ odd,} \end{cases} \end{cases} \quad (2.2)$$

Definition 2.2. A recursive relation which yields the Boubaker polynomials is:

$$\begin{cases} B_m(X) = XB_{m-1}(X) - B_{m-2}(X), & \text{for } m > 2, \\ B_2(X) = X^2 + 2, \\ B_1(X) = X, \\ B_0(X) = 1. \end{cases} \quad (2.3)$$

3. Riordan matrices of the Boubaker polynomials

In this section, we will present a Riordan matrices analysis of the Boubaker polynomials. The notations and the results of Luzon (2010a, 2008, 2009, 2010b) will be used extensively. We start with the following relation (demonstrated on page 25 in Luzon and Moron (2010b)):

$$B_n(x) = U_n\left(\frac{x}{2}\right) + 3U_{n-2}\left(\frac{x}{2}\right), \quad \text{for } n \geq 2, \quad (3.1)$$

where U_n denote the Chebyshev polynomials of the second kind. Then:

$$B_{2m}(x) = U_{2m}\left(\frac{x}{2}\right) + 3U_{2m-2}\left(\frac{x}{2}\right) = 2 \sum_{k=0}^m \tilde{T}_{2k}\left(\frac{x}{2}\right) + 6 \sum_{k=0}^{m-1} \tilde{T}_{2k}\left(\frac{x}{2}\right) \quad (3.2)$$

$$= 8 \sum_{k=0}^{m-1} \tilde{T}_{2k}\left(\frac{x}{2}\right) + 2\tilde{T}_{2m}\left(\frac{x}{2}\right) = 4 + 8 \sum_{k=0}^{m-1} \tilde{T}_{2k}\left(\frac{x}{2}\right) + 2T_{2m}\left(\frac{x}{2}\right), \quad (3.3)$$

where T_n denote the Chebyshev polynomials of the first kind. In a similar way:

$$B_{2m+1}(x) = 8 \sum_{k=0}^{m-1} \tilde{T}_{2k+1}\left(\frac{x}{2}\right) + 2\tilde{T}_{2m+1}\left(\frac{x}{2}\right) = 8 \sum_{k=0}^{m-1} T_{2k+1}\left(\frac{x}{2}\right) + 2T_{2m+1}\left(\frac{x}{2}\right) \quad (3.4)$$

$$= 8 \sum_{k=0}^{m-1} \tilde{T}_{2k}\left(\frac{x}{2}\right) + 2\tilde{T}_{2m+1}\left(\frac{x}{2}\right), \quad (3.5)$$

so:

$$\begin{aligned} B_{2m}(2 \cos t) &= 4 + 8 \sum_{k=1}^{m-1} T_{2k}(\cos t) + 2T_{2m}(\cos t) \\ &= 4 + 8 \sum_{k=1}^{m-1} \cos(2kt) + 2 \cos(2mt), \end{aligned} \quad (3.6)$$

$$B_{2m+1}(2 \cos t) = 8 \sum_{k=1}^{m-1} \cos((2k+1)t) + 2 \cos((2m+1)t). \quad (3.7)$$

Now, consider another new polynomial class \widehat{B}_n defined by:

$$\widehat{B}_n(2 \cos t) = \frac{B_n(2 \cos t) - 2T_n(\cos t)}{4}, \quad n > 1, \quad (3.8)$$

or:

$$\begin{cases} \widehat{B}_n(x) = \frac{B_n(x) - 2T_n\left(\frac{x}{2}\right)}{4}, \\ x = 2 \cos t. \end{cases} \quad (3.9)$$

So using Eqs. (3.8) and (3.9) we get:

$$\widehat{B}_{2m}(x) = \frac{B_{2m}(x) - 2T_{2m}\left(\frac{x}{2}\right)}{4} = 1 + 2 \sum_{k=0}^{m-1} \tilde{T}_{2k}\left(\frac{x}{2}\right), \quad (3.10)$$

$$\widehat{B}_{2m+1}(x) = \frac{B_{2m+1}(x) - 2T_{2m+1}\left(\frac{x}{2}\right)}{4} = 2 \sum_{k=0}^{m-1} \tilde{T}_{2k}\left(\frac{x}{2}\right). \quad (3.11)$$

In order to obtain a generating function and to make a polynomial sequence (i.e. the degree is the subindex) we consider

$$\widetilde{B}_n(x) = \widehat{B}_{n-2}(x).$$

So, symbolically:

$$\begin{bmatrix} \widetilde{B}_0(x) \\ \widetilde{B}_1(x) \\ \widetilde{B}_2(x) \\ \widetilde{B}_3(x) \\ \widetilde{B}_4(x) \\ \widetilde{B}_5(x) \\ \vdots \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 2 & 0 & 0 & 0 \\ 2 & 0 & 2 & 0 & 2 & 0 & 0 \\ 0 & 2 & 0 & 2 & 0 & 2 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \tilde{T}_0(x) \\ \tilde{T}_1(x) \\ \tilde{T}_2(x) \\ \tilde{T}_3(x) \\ \tilde{T}_4(x) \\ \tilde{T}_5(x) \\ \vdots \end{bmatrix}. \quad (3.12)$$

We can write this in terms of Riordan matrices in the next way:

$$\sum_{n \geq 0} \widetilde{B}_n(t) = T\left(\frac{2}{1-x^2}|1\right) T\left(\frac{1-x^2}{4}|\frac{1+x^2}{2}\right) T(2|2)\left(\frac{1}{1-tx}\right), \quad (3.13)$$

or:

$$\sum_{n \geq 0} \widetilde{B}_n(t)x^n = T(1|1+x^2)\left(\frac{1}{1-tx}\right). \quad (3.14)$$

In fact we have the Riordan matrix:

$$T(1|1+x^2), \quad (3.15)$$

which is:

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & -3 & 0 & 1 & 0 & 0 \\ 0 & 3 & 0 & -4 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}. \quad (3.16)$$

Hence, the few first $\tilde{B}_n(x)$ are:

$$\begin{cases} \tilde{B}_0(x) = 1, \\ \tilde{B}_1(x) = x, \\ \tilde{B}_2(x) = x^2 - 1, \\ \tilde{B}_3(x) = x^3 - 2x, \\ \tilde{B}_4(x) = x^4 - 3x^2 + 1, \\ \tilde{B}_5(x) = x^5 - 4x^3 + 3x, \end{cases} \quad (3.17)$$

with the recurrence (3.18).

$$\tilde{B}_n(x) = x\tilde{B}_{n-1}(x) - \tilde{B}_{n-2}(x), \quad n \geq 2. \quad (3.18)$$

Note that this recurrence is the same as that for the Boubaker polynomials but with different initial conditions. In fact the relation between both families of polynomials is given by

$$T(1 + 3x^2|1 + x^2) = T(1 + 3x^2|1)T(1|1 + x^2). \quad (3.19)$$

Then, finally:

$$B_n(x) = x\tilde{B}_{n-1}(x) + 3\tilde{B}_{n-2}(x), \quad n \geq 2. \quad (3.20)$$

4. Fermat-linked expressions

Using inversion of Riordan matrices we can get $\tilde{B}_n(x)$ each as combinations of Boubaker polynomials.

Remark 4.1. Comparing the recurrence (3.20) with the one of the Chebyshev polynomials of the second kind, we can obtain an explicit expression of the new polynomials defined by (3.8) and (3.9)

$$B_n(x) = \frac{\sin((n+1)t)}{\sin t}, \quad x = 2 \cos t, \quad n = 0, 1, 2, \dots \quad (4.1)$$

In other word, the new polynomial is the scaled Chebyshev polynomial $U_n(x)$ of the second kind, since the relation between the two polynomials is related as:

$$B_n(2x) = U_n(x), \quad n = 0, 1, 2, \dots \quad (4.2)$$

Remark 4.2. By using (4.1) or (4.2), we can obtain some other relations. In fact the Fermat polynomials are obtained by setting $p(x) = 3x$ and $q(x) = -2$ in the Lucas polynomial sequence, defined by (4.3).

$$F_n(x) = p(x)F_{n-1}(x) + q(x)F_{n-2}(x). \quad (4.3)$$

As Luzon (2010a), Luzon and Moron (2008, 2009, 2010b) demonstrated, through the associated Riordan matrix:

$$\begin{bmatrix} \frac{1}{3} \\ 0 & 1 \\ 0 & 0 & 3 \\ 0 & -2 & 0 & 9 \\ 0 & 0 & -12 & 0 & 27 \\ 0 & 4 & 0 & -54 & 0 & 81 \\ 0 & 0 & 36 & 0 & -216 & 0 & 243 \\ 0 & -8 & 0 & 216 & 0 & -810 & 0 & 729 \\ \vdots & \ddots \end{bmatrix}, \quad (4.4)$$

that

$$\begin{cases} F_1(x) = 1, \\ F_2(x) = 3x, \\ F_3(x) = 9x^2 - 2, \\ F_4(x) = 27x^3 - 12x, \\ \dots \quad \dots \quad \dots \end{cases} \quad (4.5)$$

and

$$F_x(x) = (\sqrt{2})^n U_n\left(\frac{3x}{2\sqrt{2}}\right). \quad (4.6)$$

Theorem 4.3. Let $(R, +, \cdot)$ be a commutative ring, $(D, +, \cdot)$ be an integral domain such that D is a subring of R whose zero is 0_D and whose unity is 1_D , $X \in R$ be transcendental over D , $D[X]$ be the ring of polynomials forms in X over D , and finally denote Boubaker polynomials and Fermat polynomials as $B_n(x)$ and $F_n(x)$, respectively, as polynomials contained in $D[X]$, then:

$$B_n(x) = \frac{1}{(\sqrt{2})^n} F_n\left(\frac{2\sqrt{2}x}{3}\right) + \frac{1}{(\sqrt{2})^{n-2}} F_{n-2}\left(\frac{2\sqrt{2}x}{3}\right); \quad n = 0, 1, 2, \dots \quad (4.7)$$

Proof. Riordan matrices for Boubaker polynomials and Fermat polynomials (see Luzon, 2010a; Luzon and Moron, 2008, 2009, 2010b) are respectively:

$$\sum_{n=0}^{+\infty} B_n(x) t^n = (1 + 3x^2|1 + x^2) \left(\frac{1}{1 - xt} \right), \quad \sum_{n=0}^{+\infty} F_n(x) t^n = \left(\frac{1}{3} \mid \frac{1+x^2}{3} \right). \quad (4.8)$$

Let's expand the inverse Riordan arrays:

$$T(1 + 3x^2|1 + x^2) = T(1 + 3x^2|1) T\left(\frac{1}{2} \mid \frac{1+x^2}{2}\right) T(2|2), \quad (4.9)$$

which gives

$$T(1 + 3x^2|1 + x^2) = T(1 + 3x^2|1) T(1|\sqrt{2}) T\left(\frac{1}{3} \mid \frac{1+x^2}{3}\right) T\left(3 \mid \frac{3}{\sqrt{2}}\right). \quad (4.10)$$

By identifying Riordan matrix for Fermat polynomials in the right term of Eq. (4.10), the desired equality holds. \square

Expressions (4.2) and (4.7) are very useful for developing the already proposed Boubaker polynomials expansion scheme (BPES).

5. Conclusion

The Boubaker polynomials have been investigated. Using a Riordan matrices (Shapiro et al., 1991) analysis, a sequence of relations outlining the relations with Chebyshev and Fermat polynomials has been obtained as guides to further studies (Yildirim and Özış, 2009, 2010). The obtained expression are a meaningful supplement to recent applied physics studies (Ben Mahmoud, 2009; Lazzez et al., 2009; Fridjine et al., 2009; Khélia et al., 2009; Ben Mahmoud and Amlouk, 2009; Dada et al., 2009; Rahmanov, xxxx; Rahmanov, xxxx; Tabatabaei et al., 2009; Fridjine and Amlouk, 2009; Belhadj et al., 2009; Belhadj et al., 2009; Zhang and Li, 2009) using the Boubaker polynomials expansion scheme (BPES).

References

- Alvarez, G., Senb, C., Furukawac, N., Motomed, Y., Dagottee, E., 2005. The truncated polynomial expansion Monte Carlo method for fermion systems coupled to classical fields: a model independent implementation. *Comput. Phys. Commun.* 168, 32–45.
- Awojoyogbe, O.B., Boubaker, K., 2009. A solution to bloch NMR flow equations for the analysis of homodynamic functions of blood flow system using m -Boubaker polynomials. *J. Current Appl. Phys.* 9, 278–283.
- Bagula, R.L., Adamson, G., 2008. Triangle of coefficients of Recursive Polynomials for Boubaker polynomials. *Encycloped. Integer Seq.* A, 137276.
- Belhadj, A., Bessrour, J., Bouhafs, M., Barrallier, L., 2009. Experimental and theoretical cooling velocity profile inside laser welded metals using keyhole approximation and Boubaker polynomials expansion. *J. Thermal Anal. Calorimetry* 97, 911–920.
- Belhadj, A., Onyango, O., Rozibaeva, N., 2009. Boubaker polynomials expansion scheme-related heat transfer investigation inside keyhole model. *J. Thermophys. Heat Trans.* 23, 639–642.
- Bender, C.M., Dunne, G.V., 1988. Polynomials and operator orderings. *J. Math. Phys.* 29, 1727–1731.
- Ben Mahmoud, B.K., 2009. Temperature 3D profiling in cryogenic cylindrical devices using Boubaker polynomials expansion scheme (BPES). *Cryogenics* 49, 217–220.
- Ben Mahmoud, K.B., Amlouk, M., 2009. The 3D Amlouk–Boubaker expansivity-energy gap-Vickers hardness abacus: a new tool for optimizing semiconductor thin film materials. *Mater. Lett.* 63 (12), 991–994.
- Boubaker, K., 2007. On modifed Boubaker polynomials: some differential and analytical properties of the new polynomial issued from an attempt for solving bi-varied heat equation. *Trends Appl. Sci. Res.* 2 (6), 540–544.
- Boubaker, K., 2008. The Boubaker polynomials, a new function class for solving bi-varied second order differential equations. *F. E. J. Appl. Math.* 31, 299–320.
- Chaouachi, A., Boubaker, K., Amlouk, M., Bouzouita, H., 2007. Enhancement of pyrolysis spray disposal performance using thermal time-response to precursor uniform deposition. *Eur. Phys. J. Appl. Phys.* 37, 105–109.
- Choi, S.K., Grandhi, R.V., Canfeld, R.A., Pettit, C.L., 2004. Polynomial chaos expansion with latin hypercube sampling for estimating response variability. *Amer. Inst. Aeronaut. Astronaut. J.* 42, 1191–1198.
- Dada, M., Awojoyogbe, O.B., Boubaker, K., 2009. Heat transfer spray model: an improved theoretical thermal time-response to uniform layers deposit using Bessel and Boubaker polynomials. *Current Appl. Phys.* 9 (3), 622–624.
- Fridjine, S., Amlouk, M., 2009. A new parameter: an ABACUS for optimizig functional materials using the Boubaker polynomials expansion scheme. *Modern Phys. Lett. B* 23, 2179–2182.
- Fridjine, S., Ben Mahmoud, K.B., Amlouk, M., Bouhafs, M., 2009. A study of sulfur/selenium substitution effects on physical and mechanical properties of vacuum-grown $ZnS1?xSex$ compounds using Boubaker polynomials expansion scheme (BPES). *J. Alloys Compounds* 479, 457–461.
- Ghanouchi, J., Labiad, H., Boubaker, K., 2008. An attempt to solve the heat transfert equation in a model of pyrolysis spray using $4q$ -order m -Boubaker polynomials. *Int. J. Heat Technol.* 26, 49–53.
- Ghrib, T., Boubaker, K., Bouhafs, M., 2008. Investigation of thermal diffusivity-microhardness correlation extended to surface-nitrured steel using Boubaker polynomials expansion. *Modern Phys. Lett. B* 22, 2893–2907.
- Guertz, B.J., van Buuren, R., Lu, H., 2000. Application of polynomial preconditioners to conservation laws Application of polynomial preconditioners to conservation laws. *J. Eng. Math.* 38, 403–426.
- Khélia, C., Boubaker, K., Ben Nasrallah, T., Amlouk, M., Belgacem, S., 2009. Morphological and thermal properties of β -SnS₂ sprayed thin films using Boubaker polynomials expansion. *J. Alloys Compounds* 477, 461–467.
- Koelink, H.T., 1994. The addition formula for continuous q -Legendre polynomials and associated spherical elements on the SU(2) quantum group related to Askey–Wilson polynomials. *SIAM J. Math. Anal.* 25 (1), 197–217.
- Labiad, H., Boubaker, K., 2007. A Sturm–Liouville shaped characteristic differential equation as a guide to establish a quasi-polynomial expression to the Boubaker polynomials. *Differ. Eqn. Control Process. Electron. J.* 2, 117–133.
- Labiad, H., Dada, M., Awojoyogbe, O.B., Ben Mahmoud, K.B., Bannour, A., 2008. Establishment of an ordinary generating function and a Christofel–Darboux type first-order differential equation for the heat equation related Boubaker–Turki polynomials. *J. Differ. Eqns.* 1, 51–66.
- Lazzez, S., Ben Mahmoud, K.B., Abroug, S., Saadallah, F., Amlouk, M., 2009. A Boubaker polynomials expansion scheme (BPES)-related protocol for measuring sprayed thin films thermal characteristics. *Current Appl. Phys.* 9, 1129–1133.
- Luzon, A., 2010a. Iterative processes related to Riordan arrays: the reciprocation and the inversion of power series. *Discrete Math.* 310, 3607–3618.
- Luzon, A., Moron, M.A., 2008. Ultrametrics, Banach's fixed point theorem and the Riordan group. *Discrete Appl. Math.* 156, 2620–2635.
- Luzon, A., Moron, M.A., 2009. Riordan matrices in the reciprocation of quadratic polynomials. *Linear Alg. Appl.* 430, 2254–2270.
- Luzon, A., Moron, M.A., 2010b. Recurrence relations for polynomial sequences via Riordan matrices. *Linear Alg. Appl.* 433, 1422–1446.
- Okada, R., Nakata, N., Spencer, B.F., Kasai, K., Saang, B.K., 2006. Rational polynomial approximation modeling for analysis of structures with VE dampers. *J. Earthquake Eng.* 10, 97–125.
- Philippou, A.N., Georgiou, C., 1989. Convolutions of Fibonacci-type polynomials of order k and the negative binomial distributions of the same order. *Fibonacci Quart.* 27, 209–216.
- Rahmanov, H. Triangle read by rows: row n gives coefficients of Boubaker polynomial $B_n(x)$, calculated for $X = 2\cos(t)$, centered by adding $-2\cos(nt)$, then divided by 4, in order of decreasing exponents. *Encycloped. Integer Seq.* A 160242.
- Rahmanov, H. Triangle read by rows: row n gives values of the 4q-28 Boubaker polynomials $B_{4q}(X)$ (named after Boubaker Boubaker (1897–1966)), calculated for $X = 1$ (or -1). *Encycloped. Integer Seq.* A 162180.
- Shapiro, L.W., Getu, S., Woan, W.J., Woodson, L.C., 1991. The Riordan Group. *Discrete Appl. Math.* 34, 229–239.

- Slama, S., Bessrour, J., Boubaker, K., Bouhafs, M., 2008a. Investigation of A3 point maximal front spatial evolution during resistance spot welding using $4q$ -Boubaker polynomial sequence. In: Proceedings of COTUME, pp. 79–80.
- Slama, S., Bessrour, J., Boubaker, K., Bouhafs, M., 2008b. A dynamical model for investigation of A3 point maximal spatial evolution during resistance spot welding using Boubaker polynomials. *Eur. Phys. J. Appl. Phys.* 44, 317–322.
- Slama, S., Bouhafs, M., Ben Mahmoud, K.B., 2008c. A Boubaker polynomials solution to heat equation for monitoring A3 point evolution during resistance spot welding. *Int. J. Heat Technol.* 26 (2), 141–146.
- Sloan, I.H., Womersley, R.S., 2002. Good approximation on the sphere, with application to geodesy and the scattering of sound. *J. Comput. Appl. Math.* 149, 227–237.
- Sloane, N.J.A., 2008. Triangle read by rows of coefficients of Boubaker polynomial $B_n(x)$ in order of decreasing exponents. Encycloped. Integer Seq., A138034.
- Tabatabaei, S., Zhao, T., Awojoyogbe, O., Moses, F., 2009. Cut-off cooling velocity profiling inside a keyhole model using the Boubaker polynomials expansion scheme. *Int. J. Heat Mass Transfer* 45, 1247–1255.
- Yildirim, A., 2010. He's homotopy perturbation method for nonlinear differential-difference equations. *Int. J. Comput. Math.* 87, 992–996.
- Yildirim, A., Özış, T., 2009. Solutions of Singular IVPs of Lane-Emden type by the variational iteration method. *Nonlinear Anal. Series A: Theor. Methods & Appl.* 70, 2480–2484.
- Zhang, D.H., Li, F.W., 2009. A Boubaker polynomials expansion scheme BPES-related analytical solution to Williams–Brinkmann stagnation point flow equation at a blunt body. *Int. J. Appl. Phys. Lett.* 2, 25–30.
- Zhao, T.G., Ben Mahmoud, B.K., Toumi, M.A., Faromika, O.P., Dada, M., Awojoyogbe, O.B., Magnuson, J., Lin, F., 2009. Some new properties of the applied-physics related Boubaker polynomials. *Differ. Eqns. Control Process.* 1, 7–19.
- Zhao, T.G., Wang, Y.X., Ben Mahmoud, B.K., 2010. Limit and uniqueness of the Boubaker–Zhao polynomials single imaginary root sequence. *Int. J. Math. Comput.* 1(8), 13–16.